

# The ZX-calculus is incomplete for quantum mechanics

Christian Schröder de Witt <sup>1</sup>   Vladimir Zamdzhiev <sup>2</sup>

<sup>1</sup>Scherenbergstr. 22, 10439 Berlin

<sup>2</sup>Department of Computer Science,  
University of Oxford

5 June 2014

# ZX-calculus

- Introduced by Coecke and Duncan in 2008

# ZX-calculus

- Introduced by Coecke and Duncan in 2008
- Diagrammatic logical calculus for studying quantum information processing

# ZX-calculus

- Introduced by Coecke and Duncan in 2008
- Diagrammatic logical calculus for studying quantum information processing
- Can be used as an alternative to traditional Hilbert space formalism

# ZX-calculus

- Introduced by Coecke and Duncan in 2008
- Diagrammatic logical calculus for studying quantum information processing
- Can be used as an alternative to traditional Hilbert space formalism
- Has been used to study:
  - Quantum algorithms
  - Quantum security protocols
  - Quantum error-correcting codes
  - and other problems involving quantum information

## Atomic Diagrams (1)

$$\left[ \begin{array}{|c|} \hline | \\ \hline \end{array} \right] = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I \quad \left[ \begin{array}{|c|} \hline \diagdown \\ \diagup \\ \hline \end{array} \right] = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \sigma$$

$$\left[ \begin{array}{|c|} \hline \cap \\ \hline \end{array} \right] = \langle 00| + \langle 11| \quad \left[ \begin{array}{|c|} \hline \cup \\ \hline \end{array} \right] = |00\rangle + |11\rangle$$

$$\left[ \begin{array}{|c|} \hline \text{H} \\ \hline \end{array} \right] = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = H$$

# Atomic Diagrams (2)

$$\left[ \begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \text{0} \\ \vdots \\ \vdots \\ \vdots \end{array} \right] = \begin{cases} |0^m\rangle & \mapsto |0^n\rangle \\ |1^m\rangle & \mapsto e^{i\alpha} |1^n\rangle \\ \text{rest} & \mapsto 0 \end{cases}$$

$$\left[ \begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \alpha \\ \vdots \\ \vdots \\ \vdots \end{array} \right] = \begin{cases} |+\!^m\rangle & \mapsto |+\!^n\rangle \\ |-\!^m\rangle & \mapsto e^{i\alpha} |-\!^n\rangle \\ \text{rest} & \mapsto 0 \end{cases}$$

where  $\alpha \in [0, 2\pi)$

## Compound Diagrams

$$\left[ \begin{array}{c} \dots \\ \text{---} \\ \text{---} \\ \Psi_1 \\ \text{---} \\ \text{---} \\ \dots \end{array} \right] = D_1 \quad \text{and} \quad \left[ \begin{array}{c} \dots \\ \text{---} \\ \text{---} \\ \Psi_2 \\ \text{---} \\ \text{---} \\ \dots \end{array} \right] = D_2$$

then

$$\left[ \begin{array}{cc} \dots & \dots \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \Psi_1 & \Psi_2 \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \dots & \dots \end{array} \right] = D_1 \otimes D_2$$

and

$$\left[ \begin{array}{c} \dots \\ \text{---} \\ \text{---} \\ \Psi_1 \\ \text{---} \\ \text{---} \\ \dots \\ \text{---} \\ \text{---} \\ \Psi_2 \\ \text{---} \\ \text{---} \\ \dots \end{array} \right] = D_1 \circ D_2$$



## Examples

$$\llbracket \text{red circle} \rrbracket = \sqrt{2} |0\rangle$$

$$\llbracket \text{red circle with } \pi \rrbracket = \sqrt{2} |1\rangle$$

$$\llbracket \text{green circle} \rrbracket = \sqrt{2} |+\rangle$$

$$\llbracket \text{green circle with } \pi \rrbracket = \sqrt{2} |-\rangle$$

$$\llbracket \text{green circle with } \pi \text{ and wire} \rrbracket = Z$$


$$\llbracket \text{red circle with } \pi \text{ and wire} \rrbracket = X$$

$$\llbracket \text{green circle with wire connected to red circle} \rrbracket = \text{CNOT}$$



## Axioms (1)

*"Only the topology matters"*


## Example




## Axioms (2)




# Axioms (3)




# Axioms (4)




# Axioms (5)



# Axioms(6)



# Example derivation





# Soundness, Completeness and Universality results

- The ZX-calculus is sound

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal
  - $\forall U : \mathcal{Q}^n \rightarrow \mathcal{Q}^m, \exists D. \llbracket D \rrbracket = U$

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal
  - $\forall U : \mathcal{Q}^n \rightarrow \mathcal{Q}^m, \exists D. \llbracket D \rrbracket = U$
- Completeness of ZX (with the Euler decomposition of the Hadamard gate rule) was unknown.

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal
  - $\forall U : \mathcal{Q}^n \rightarrow \mathcal{Q}^m, \exists D. \llbracket D \rrbracket = U$
- Completeness of ZX (with the Euler decomposition of the Hadamard gate rule) was unknown.
  - $\llbracket D_1 \rrbracket = \llbracket D_2 \rrbracket \implies ZX \vdash D_1 = D_2$

# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal
  - $\forall U : \mathcal{Q}^n \rightarrow \mathcal{Q}^m, \exists D. \llbracket D \rrbracket = U$
- Completeness of ZX (with the Euler decomposition of the Hadamard gate rule) was unknown.
  - $\llbracket D_1 \rrbracket = \llbracket D_2 \rrbracket \implies ZX \vdash D_1 = D_2$
- The ZX-calculus is complete for stabilizer quantum mechanics


# Soundness, Completeness and Universality results

- The ZX-calculus is sound
  - $ZX \vdash D_1 = D_2 \implies \llbracket D_1 \rrbracket = e^{i\phi} \llbracket D_2 \rrbracket$
- The ZX-calculus is universal
  - $\forall U : \mathcal{Q}^n \rightarrow \mathcal{Q}^m, \exists D. \llbracket D \rrbracket = U$
- Completeness of ZX (with the Euler decomposition of the Hadamard gate rule) was unknown.
  - $\llbracket D_1 \rrbracket = \llbracket D_2 \rrbracket \implies ZX \vdash D_1 = D_2$
- The ZX-calculus is complete for stabilizer quantum mechanics
  - If  $D_1$  and  $D_2$  are ZX-SQM diagrams and  $\llbracket D_1 \rrbracket = \llbracket D_2 \rrbracket$  then  $ZX \vdash D_1 = D_2$



# Euler Decomposition


Recall that for single-qubit unitary maps:



where  $\alpha_i, \beta_i, \gamma_i, \phi_i \in [0, 2\pi)$

## Alternative Models


Consider the following models:



$$[[\cdot]]_k := [[\cdot]] , \text{ otherwise}$$

These models are sound when  $k = 4p + 1$  for  $p \in \mathbb{Z}$ .

# Counter-example diagrams



# Counter-example diagrams (cont.)

where

$$\alpha := -\arccos\left(\frac{5}{2\sqrt{13}}\right) \approx 0.2561\pi$$

$$\beta := -2\arcsin\left(\frac{\sqrt{3}}{4}\right) \approx -0.2851\pi$$

$$\phi := \arcsin\left(\frac{\sqrt{3}}{4}\right) - \alpha \approx 0.3987\pi$$

# Incomplete

We have:

$$\llbracket D_1 \rrbracket = \llbracket D_2 \rrbracket$$

but for any  $\lambda \in \mathbb{C}$  :

$$\llbracket D_1 \rrbracket_{-3} \neq \lambda \llbracket D_2 \rrbracket_{-3}$$

However,  $\llbracket \cdot \rrbracket_{-3}$  is a sound model of ZX, so

$$ZX \not\vdash D_1 = D_2$$

and therefore the ZX-calculus is incomplete for quantum mechanics.

## ZX is incomplete, what next?


- Nothing special about counter-example, others easily doable using the same approach

## ZX is incomplete, what next?

- Nothing special about counter-example, others easily doable using the same approach
- Color-swap rule might be needed for completeness

# ZX is incomplete, what next?

- Nothing special about counter-example, others easily doable using the same approach
- Color-swap rule might be needed for completeness




where

$$\begin{array}{lll} \alpha_2 := f_1(\alpha_1, \beta_1, \gamma_1) & ; & f_1 = ? \\ \beta_2 := f_2(\alpha_1, \beta_1, \gamma_1) & ; & f_2 = ? \\ \gamma_2 := f_3(\alpha_1, \beta_1, \gamma_1) & ; & f_3 = ? \end{array}$$



# Even then, more challenges



# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness

## Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):

# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds

# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds
  - calculus is not even approximately universal

# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds
  - calculus is not even approximately universal
- If diagram angles are of the form  $\frac{k\pi}{4}$  (Clifford+T):

# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds
  - calculus is not even approximately universal
- If diagram angles are of the form  $\frac{k\pi}{4}$  (Clifford+T):
  - calculus is approximately universal

# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds
  - calculus is not even approximately universal
- If diagram angles are of the form  $\frac{k\pi}{4}$  (Clifford+T):
  - calculus is approximately universal
  - calculus is complete for line-graphs (next talk)



# Approximate completeness instead?

- Restrict generators, drop universality and work towards approximate completeness
- If diagram angles are of the form  $\frac{k\pi}{2}$  (Stabilizer QM):
  - completeness holds
  - calculus is not even approximately universal
- If diagram angles are of the form  $\frac{k\pi}{4}$  (Clifford+T):
  - calculus is approximately universal
  - calculus is complete for line-graphs (next talk)
  - completeness is unknown